arXiv:1208.1965v1 [quant-ph] 9 Aug 2012

Quantum nonlocal boxes and nonlocal box swapping

Adán Cabello
Departamento de Física Aplicada II, Universidad de Sevilla, E-41012 Sevilla, Spain

(Dated: August 10, 2012)

Nonlocal boxes (NBs) are theoretical objects that violate a bipartite Bell inequality as much as the no-signaling principle allows. For most bipartite scenarios, NBs are physically impossible to prepare. We show that NBs are physically feasible if the two parties have 3 measurements with 4 outputs. Moreover, we show that, in this case, there is a quantum protocol for NB swapping which allows the parties to locally prepare mixtures of NBs using systems that never interacted. A stronger form of bipartite quantum nonlocality emerges in this scenario: systems that never interacted are as nonlocal as allowed by no-signaling.

Introduction.—The violation of the Clauser-Horne-Shimony-Holt (CHSH) Bell inequality [1, 2] has been experimentally observed (up to loopholes [3]) many times [4–7] and has a number of applications [8–10]. However, although counterintuitive, this nonlocality is not as strong as it might be. On one hand, it occurs for systems that have interacted in the past, rather than for systems that have never interacted. On the other hand, the maximum quantum violation [11] is not as large as the one allowed by the no-signaling principle that prevents causal influences from propagating faster-than-light [12]; therefore hypothetical postquantum resources could give higher violations. Here we show that, under the appropriate conditions, both "limitations" vanish and, as a result, a stronger form of quantum nonlocality emerges.

Quantum nonlocal boxes.—Nonlocal boxes (NBs) are theoretical objects shared by two parties, Alice and Bob, that: (I) allow them to violate a Bell inequality as much as is allowed by the no-signaling principle, and (II) enable each party to predict with certainty the outcome of the other party. NBs were introduced to show that the maximum quantum violation of the CHSH inequality is not as large as allowed by the no-signaling principle [12]. For the CHSH and other Bell inequalities, NBs are rendered impossible by some physical principles [13–15].

A complete set of NBs consists of NBs of different classes which can be combined to produce any desired correlations between Alice and Bob, ranging from totally uncorrelated results (i.e., each party cannot infer anything about the other party's result) to correlations maximally violating a Bell inequality.

Our first result is the following: If Alice and Bob can choose between 3 experiments with 4 outcomes, then a third party, Eve, can prepare a complete set of NBs.


A_0 A_1 A_2
-- -+ +- ++
Alice
B_0 B_1 B_2
-- -+ +- ++
Bob
t
(ii)
(iii)
1, 3
2, 4
(i)
x
y
ϕ^+_{12} ϕ^-_{12} ψ^+_{12} ψ^-_{12}
ϕ^+_{34} ϕ^-_{34} ψ^+_{34} ψ^-_{34}
Eve

FIG. 1: **Quantum nonlocal boxes.** Space-time diagram of a run of an experiment in which Eve produces a NB. In (i), Eve prepares 4 qubits in a product of Bell states, e.g., qubit 1 and 2 in the state ϕ^-_{12} and qubits 3 and 4 in the state ϕ^+_{34}. In (ii), Alice presses a button, e.g., A_2 (leading to observable A_2 being measured on qubits 1 and 3) and records the result (e.g., $-+$). In (iii), spacelike separated form (ii), Bob presses a button, e.g., B_1 (leading to observable B_1 being measured on qubits 2 and 4) and records the result (e.g., $--$). Eve previously set observable A_i and B_i to be those defined in (3) and (5), respectively.

We prove it by constructing an explicit example which is easy to follow with the aid of black boxes with buttons; see Fig. 1. For each run of the experiment, Alice and Bob randomly choose one out of 3 buttons, representing the experiments they can perform on a system prepared by Eve. Alice's buttons are labeled A_0, A_1, A_2, and Bob's B_0, B_1, B_2. Every time one button is pressed, one light flashes, indicating the corresponding result. There are 4 lights, and we label each of them with two bits: $--$ [meaning $(-1,-1)$], $-+$, $+-$, and $++$. There is spacelike separation between the event in which Alice (Bob) presses the button and the event in which Bob (Alice) records the result, so that there cannot be causal influences between them.

Consider the following 16 Bell inequalities:

$$\beta_i \overset{\text{LHV}}{\leq} 7 \overset{\text{NS}}{\leq} 9, \tag{1}$$

where $i = 1, \ldots, 16$, $\overset{\text{LHV}}{\leq} 7$ indicates that, for any local hidden variable theory, β_i is upper bounded by 7, $\overset{\text{NS}}{\leq} 9$ indicates that, for any theory satisfying no-signaling, β_i is upper bounded by 9, and

$$\begin{aligned}\beta_i =& s_{00}\langle A_0^{10}B_0^{10}\rangle + s_{01}\langle A_0^{01}B_1^{10}\rangle + s_{02}\langle A_0^{11}B_2^{10}\rangle \\ &+ s_{10}\langle A_1^{10}B_0^{01}\rangle + s_{11}\langle A_1^{01}B_1^{01}\rangle + s_{12}\langle A_1^{11}B_2^{01}\rangle \\ &+ s_{20}\langle A_2^{10}B_0^{11}\rangle + s_{21}\langle A_2^{01}B_1^{11}\rangle + s_{22}\langle A_2^{11}B_2^{11}\rangle,\end{aligned} \tag{2}$$

where $\langle A_1^{10}B_0^{01}\rangle$ is the mean value of the product of the first bit (-1 or 1) of the result of measuring A_1, multiplied by the second bit of the result of B_0, and $\langle A_2^{01}B_1^{11}\rangle$ is the mean value of the product of the second bit of the result of A_2 times the first bit of the result of B_1 times the second bit of the result of B_1. For each β_i, the values of $s_{00}, \ldots, s_{22}$ are given in Table I. It can be checked (e.g., using `porta` [16]), that all these 16 inequalities are tight (i.e., facets of the bipartite 4-measurement, 3-outcome local polytope [17]).

TABLE I: Coefficients s_j in (2).

	s_{00}	s_{01}	s_{02}	s_{10}	s_{11}	s_{12}	s_{20}	s_{21}	s_{22}
$\beta_{1,16}$	±1	±1	1	±1	±1	1	1	1	-1
$\beta_{2,15}$	±1	±1	1	∓1	±1	-1	-1	1	1
$\beta_{3,14}$	±1	∓1	-1	±1	±1	1	1	-1	1
$\beta_{4,13}$	±1	∓1	-1	∓1	±1	-1	-1	-1	-1
$\beta_{5,12}$	±1	±1	1	±1	∓1	-1	1	-1	1
$\beta_{6,11}$	±1	±1	1	∓1	∓1	1	-1	-1	-1
$\beta_{7,10}$	±1	∓1	-1	±1	∓1	-1	1	1	-1
$\beta_{8,9}$	±1	∓1	-1	∓1	∓1	1	-1	1	1

Eve sets Alice's A_0, A_1, and A_2 to the following two-qubit observables:

$$\begin{aligned}A_0 =& r_{++}|00\rangle\langle 00| + r_{+-}|01\rangle\langle 01| \\ &+ r_{-+}|10\rangle\langle 10| + r_{--}|11\rangle\langle 11|,\end{aligned} \tag{3a}$$

$$\begin{aligned}A_1 =& r_{++}|++\rangle\langle ++| + r_{+-}|-+\rangle\langle -+| \\ &+ r_{-+}|+-\rangle\langle +-| + r_{--}|--\rangle\langle --|,\end{aligned} \tag{3b}$$

$$\begin{aligned}A_2 =& r_{++}|\chi^+\rangle\langle\chi^+| + r_{+-}|\chi^-\rangle\langle\chi^-| \\ &+ r_{-+}|\omega^+\rangle\langle\omega^+| + r_{--}|\omega^-\rangle\langle\omega^-|,\end{aligned} \tag{3c}$$

where $\sigma_z|0\rangle = |0\rangle$, $\sigma_z|1\rangle = -|1\rangle$, $\sigma_x|\pm\rangle = \pm|\pm\rangle$, and

$$|\chi^\pm\rangle = \frac{1}{\sqrt{2}}(|0+\rangle \pm |1-\rangle), \tag{4a}$$

$$|\omega^\pm\rangle = \frac{1}{\sqrt{2}}(|1+\rangle \pm |0-\rangle) \tag{4b}$$

are the common eigenvectors of $\sigma_z \otimes \sigma_x$ and $\sigma_x \otimes \sigma_z$.

Similarly, Eve sets Bob's B_0, B_1, and B_2 to the following two-qubit observables:

$$\begin{aligned}B_0 =& r_{++}|0+\rangle\langle 0+| + r_{+-}|0-\rangle\langle 0-| \\ &+ r_{-+}|1+\rangle\langle 1+| + r_{--}|1-\rangle\langle 1-|,\end{aligned} \tag{5a}$$

$$\begin{aligned}B_1 =& r_{++}|+0\rangle\langle +0| + r_{+-}|-0\rangle\langle -0| \\ &+ r_{-+}|+1\rangle\langle +1| + r_{--}|-1\rangle\langle -1|,\end{aligned} \tag{5b}$$

$$\begin{aligned}B_2 =& r_{++}|\phi^+\rangle\langle\phi^+| + r_{+-}|\phi^-\rangle\langle\phi^-| \\ &+ r_{-+}|\psi^+\rangle\langle\psi^+| + r_{--}|\psi^-\rangle\langle\psi^-|,\end{aligned} \tag{5c}$$

where

$$|\phi^\pm\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle), \tag{6a}$$

$$|\psi^\pm\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle) \tag{6b}$$

are the Bell states, which are the common eigenvectors of $\sigma_z \otimes \sigma_z$ and $\sigma_x \otimes \sigma_x$. A_i and B_i have possible outcomes r_{++} [representing $(+1, +1)$], r_{+-}, r_{-+}, and r_{--}.

In addition, Eve has to prepare 4 qubits (qubits 1 and 3 for Alice, and qubits 2 and 4 for Bob) in one of the following 16 4-qubit states: $\phi_{12}^+\phi_{34}^+$, $\phi_{12}^+\phi_{34}^-$, $\ldots$, $\psi_{12}^-\psi_{34}^-$. As shown in Table II, each of these states violates one of the 16 Bell inequalities $\beta_i \overset{\text{LHV}}{\leq} 7$ (for $i = 1, \ldots, 16$) up to its algebraic maximum, 9, which is also the maximum value allowed by the no-signaling principle. This occurs because, from each local experiment, one of the parties can predict with certainty the other party's result. For instance, from the result of A_0, Alice can predict with certainty Bob's result for B_0^{10}, B_1^{01}, and B_2^{10}. The completeness of this set of 16 NBs follows from the fact that the 16 products of Bell states form an orthogonal basis of the Hilbert space of dimension 16.

Nonlocal box swapping.—Suppose that Eve wants to prepare an equally weighted mixture of NBs, so that Alice and Bob's results look totally uncorrelated. Eve can do it by producing, at will, quartets of qubits in the 16 states with equal frequency. However, there is an alternative way to make Alice and Bob share an equally weighted mixture of NBs in which Eve's intervention is unnecessary.

This alternative is based on entanglement swapping between two pairs of 4-dimensional systems. For pairs of qubits, entanglement swapping [18] works as follows: Initially, two parties, Alice and Bo, share two qubits in a Bell state and two other parties, Al and Bob, share another two qubits in a Bell state. A measurement which distinguishes the 4 Bell states on Al's and Bo's qubits does not only project them into one of the 4 Bell states, but also projects the state of Alice's and Bob's qubit into one of the 4 Bell states. Indeed, there is a one-to-one correspondence between Al-Bo's final Bell state and Alice-Bob's final Bell state. Therefore, Alice and Bob end up sharing a Bell state, known by Al and Bo, without any interaction between their qubits having taken place.

TABLE II: Values of β_i for the 16 products of Bell states. Values in bold indicate the maximum violation allowed by no-signaling.

	β_1	β_2	β_3	β_4	β_5	β_6	β_7	β_8	β_9	β_{10}	β_{11}	β_{12}	β_{13}	β_{14}	β_{15}	β_{16}
$\phi^+_{12}\phi^+_{34}$	**9**	1	1	−3	1	1	1	−3	1	1	1	−3	−3	−3	−3	1
$\phi^+_{12}\phi^-_{34}$	1	**9**	−3	1	1	1	−3	1	1	1	−3	1	−3	−3	1	−3
$\phi^+_{12}\psi^+_{34}$	1	−3	**9**	1	1	−3	1	1	1	−3	1	1	−3	1	−3	−3
$\phi^+_{12}\psi^-_{34}$	−3	1	1	**9**	−3	1	1	1	−3	1	1	1	1	−3	−3	−3
$\phi^-_{12}\phi^+_{34}$	1	1	1	−3	**9**	1	1	−3	−3	−3	−3	1	1	1	1	−3
$\phi^-_{12}\phi^-_{34}$	1	1	−3	1	1	**9**	−3	1	−3	−3	1	−3	1	1	−3	1
$\phi^-_{12}\psi^+_{34}$	1	−3	1	1	1	−3	**9**	1	−3	1	−3	−3	1	−3	1	1
$\phi^-_{12}\psi^-_{34}$	−3	1	1	1	−3	1	1	**9**	1	−3	−3	−3	−3	1	1	1
$\psi^+_{12}\phi^+_{34}$	1	1	1	−3	−3	−3	−3	1	**9**	1	1	−3	1	1	1	−3
$\psi^+_{12}\phi^-_{34}$	1	1	−3	1	−3	−3	1	−3	1	**9**	−3	1	1	1	−3	1
$\psi^+_{12}\psi^+_{34}$	1	−3	1	1	−3	1	−3	−3	1	−3	**9**	1	1	−3	1	1
$\psi^+_{12}\psi^-_{34}$	−3	1	1	1	1	−3	−3	−3	−3	1	1	**9**	−3	1	1	1
$\psi^-_{12}\phi^+_{34}$	−3	−3	−3	1	1	1	1	−3	1	1	1	−3	**9**	1	1	−3
$\psi^-_{12}\phi^-_{34}$	−3	−3	1	−3	1	1	−3	1	1	1	−3	1	1	**9**	−3	1
$\psi^-_{12}\psi^+_{34}$	−3	1	−3	−3	1	−3	1	1	1	−3	1	1	1	−3	**9**	1
$\psi^-_{12}\psi^-_{34}$	1	−3	−3	−3	−3	1	1	1	−3	1	1	1	−3	1	1	**9**

The protocol for NB swapping is illustrated in Fig. 2. Initially, Alice and Bo share a NB based on a Bell state between qubits 1 and 2, another Bell state between qubits 3 and 4, and the observables A_i and B_j given by (3) and (5). Similarly, Al and Bob share a NB based on a Bell state between qubits 5 and 6, another Bell state between qubits 7 and 8, and the observables A_i and B_j. An entanglement swap occurs when a measurement distinguishing the 4 Bell states is performed on qubits 2 and 5, and a similar measurement is performed on qubits 4 and 7: Qubits 2 and 5 (4 and 7) end up in one of the 4 Bell states and qubits 1 and 6 (3 and 8) end up in the corresponding Bell state. As a consequence, a NB swap takes place and Alice and Bob end up sharing a NB.

Maximum nonlocality of systems that never interacted.—The protocol for NB swapping can be used to locally prepare equally weighted mixtures of NBs and then achieve maximum nonlocality allowed by no-signaling of systems that never interacted. The protocol is shown in Fig. 3. The process starts when two distant independent sources emit quartets of qubits in products of Bell states. In one region of space-time, Alice performs local measurements on pairs of qubits. In a causally unconnected region, Bob performs similar measurements. Notice that, so far, for any observer, the quantum state of the 4 qubits tested by Alice and Bob is maximally mixed (therefore unentangled). However, a robot, in a causally unconnected region from those in which Alice and Bob are performing the measurements, can sort the events in subsets such that each of them maximally violates one of the 16 Bell inequalities (1). For that, the robot performs measurements distinguishing the 16 products of Bell states.

Contemplating the whole experiment, we notice that we ended up with a nonlocality experiment free of the two limitations we mentioned. On one hand, the Bell test is performed on systems that never interacted. On the other hand, *by a measurement performed in a region spacelike separated from both Alice's and Bob's measurements*, the events are sorted in sets such that each set violates a Bell inequality up to its no-signaling limit.

Relation to previous results.—Entanglement swapping was used together with CHSH inequalities in the delayed entanglement swapping protocol [19]. However, there, Eve sorts Alice's and Bob's results in subsets not maximally violating a Bell inequality, and does it *after* the results are already recorded (thus Eve knows Alice's and Bob's choices).

The Bell inequality $\beta_{16} \overset{\text{LHV}}{\leq} 7$ was introduced in [20] in the context of all-versus-nothing proofs. The Bell inequality $\beta_1 \overset{\text{LHV}}{\leq} 7$ has been recently used to experimentally obtain nonlocal correlations with a very small local part [21].

Proposed experimental realization.—To experimentally prepare a complete set of quantum NBs (the experiment in Fig. 1), we can use pairs of photons entangled simultaneously in two different degrees of freedom (hyperentanglement). All observables A_i and B_i in (3) and (5) have been measured using the polarization and spatial degrees of a photon on a 2-photon system hyperentangled in both polarization and spatial degrees of freedom prepared in one of the 16 states needed [21, 22]. Since, given one of these states, any of the others can be obtained by a local unitary operation, there is no difficulty in preparing the complete set of 16 states. An experimental challenge is to achieve spatial separation between the local measure-



FIG. 2: **Nonlocal box swapping**. (a) Initially, Alice and Bo share a NB based on the observables A_i and B_j in (3) and (5), and in 4 qubits such that 1 and 2 (3 and 4) are in a Bell state. Al and Bob share a similar NB. (b) A measurement distinguishing among the 4 Bell states is performed on qubits 2 and 5, and a similar measurement is performed on qubits 4 and 7. (c) As a result, an entanglement swap occurs: Qubits 2 and 5 (4 and 7) end up in one of the 4 Bell states, and qubits 1 and 6 (3 and 8) end up in the corresponding Bell state. As a consequence, a NB swap also takes place, so Alice and Bob end up sharing a NB without any interaction between their qubits having taken place.



FIG. 3: **Maximum nonlocality of systems that never interacted**. (i) A source emits qubits 1 and 2 in the Bell state ψ^-_{12} and qubits 3 and 4 in the state ψ^-_{34}. (ii) A similar source, synchronized with (i), emits qubits 5 and 6 in the state ψ^-_{56} and qubits 7 and 8 in the state ψ^-_{78}. (iii) Alice performs a measurement A_i on qubits 1 and 3. (iv) Bob performs a measurement B_j on qubits 6 and 8. The quantum state of qubits 1, 3, 6, and 8 is maximally mixed. (v) A robot performs a measurement distinguishing the 4 Bell states on qubits 2 and 5, and a similar measurement on qubits 4 and 7, and use their results to sort the event in one of the 16 classes $\phi^+_{16}\phi^+_{38}, \ldots, \psi^-_{16}\psi^-_{38}$. The region of space-time in which the robot sorts the event is spacelike separated from those in which Alice and Bob perform the local measurements.

ments on the two photons in order to guarantee spacelike separation (in previous experiments [21, 22] the distance between Alice and Bob is less than a meter).

The experiment to demonstrate NB swapping and maximum nonlocality allowed by no-signaling of systems that have not interacted (the experiment in Fig. 3) is much more demanding. It requires double entanglement (since we need to swap the entanglement between two pairs of 4-dimensional systems), 3 mutually spacelike separated regions (Alice's, Bob's, and the robot's), and systems in which we can measure the 4-dimensional local observables A_i and B_i. One possibility is to achieve 4-qubit Bell state discrimination by using additional degrees of freedom, as proposed (for two-qubit Bell state discrimination) in [23] and demonstrated in [24, 25]. The challenge is to extend this idea to more degrees of freedom. For example, we could achieve 4-qubit Bell state discrimination in polarization and spatial degrees of freedom by using an 8-qubit system which includes time-energy entanglement [26] and also entanglement in an extra spatial degree of freedom [27]. Alternatively, other recent proposals for distinguishing the 16 states [28, 29] could be adopted.

Conclusions.—Nature offers a much stronger form of nonlocality than the one observed in the experiments of the CHSH inequality and exploited for a number of applications. Here we have shown that, if the parties can choose among a larger number of measurements and outcomes, then quantum NBs providing the maximum bipartite nonlocality allowed by the no-signaling principle become possible. In addition, for these NBs, a NB swapping protocol exists which permits to prepare equally weighted mixtures of NBs between two parties that have never interacted. The final picture is that of a much stronger form of nonlocality, in which Alice and Bob perform their local measurements on systems that have never interacted, and found that every pair is as nonlocal as allowed by no-signaling. This nonlocality can be observed experimentally with current technology and may lead to new applications in quantum information processing.

The author thanks A. Acín, C. Budroni, R. Gallego, and P. Mataloni for useful conversations, C.B. and R.G. for independently checking the tightness of the Bell inequalities (1) and (2), and A. J. López-Tarrida for his help in Figs. 1 and 3. This work was supported by Project No. FIS2011-29400 (Spain).

[1] J. S. Bell, Physics **1**, 195 (1964).
[2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969); *ibid.* **24**, 549 (1970).
[3] A. Cabello and F. Sciarrino, Phys. Rev. X **22**, 021010 (2012).
[4] A. Aspect, J. Dalibard, and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
[5] W. Tittel, J. Brendel, H. Zbinden, and N. Gisin, Phys. Rev. Lett. **81**, 3563 (1998).
[6] G. Weihs, T. Jennewein, C. Simon, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. **81**, 5039 (1998).
[7] D. N. Matsukevich, P. Maunz, D. L. Moehring, S. Olmschenk, and C. Monroe, Phys. Rev. Lett. **100**, 150404 (2008).
[8] Č. Brukner, M. Żukowski, J.-W. Pan, and A. Zeilinger, Phys. Rev. Lett. **95**, 010503 (2005).
[9] J. Barrett, L. Hardy, and A. Kent, Phys. Rev. Lett. **95**, 010503 (2005).
[10] S. Pironio *et al.*, Nature **464**, 1021 (2010).
[11] B. S. Cirel'son, Lett. Math. Phys. **4**, 93 (1980).
[12] S. Popescu and D. Rohrlich, Found. Phys. **24**, 379 (1994).
[13] W. van Dam, Ph.D. thesis, University of Oxford, 2000.
[14] M. Pawłowski, T. Paterek, D. Kaszlikowski, V. Scarani, A. Winter, and M. Żukowski, Nature (London) **461**, 1101 (2009).
[15] J. Oppenheim and S. Wehner, Science **330**, 1072 (2010).
[16] `http://www.zib.de/Optimization/Software/porta/`
[17] I. Pitowsky, *Quantum Probability-Quantum Logic* (Springer, Berlin, 1989).
[18] M. Żukowski, A. Zeilinger, M. A. Horne, and A. K. Ekert, Phys. Rev. Lett. **71**, 4287 (1993).
[19] A. Peres, J. Mod. Opt. **47**, 139 (2000).
[20] A. Cabello, Phys. Rev. Lett. **87**, 010403 (2001).
[21] L. Aolita, R. Gallego, A. Acín, A. Chiuri, G. Vallone, P. Mataloni, and A. Cabello, Phys. Rev. A **85**, 032107 (2012).
[22] T. Yang, Q. Zhang, J. Zhang, J. Yin, Z. Zhao, M. Żukowski, Z.-B. Chen, and J.-W. Pan, Phys. Rev. Lett. **95**, 240406 (2005).
[23] S. P. Walborn, S. Pádua, and C. H. Monken, Phys. Rev. A **68**, 042313 (2003).
[24] M. Barbieri, G. Vallone, P. Mataloni, and F. De Martini, Phys. Rev. A **75**, 042317 (2007).
[25] T.-C. Wei, J. T. Barreiro, and P. G. Kwiat, Phys. Rev. A **75**, 060305(R) (2007).
[26] J. T. Barreiro, N. K. Langford, N. A. Peters, and P. G. Kwiat, Phys. Rev. Lett. **95**, 260501 (2005).
[27] R. Ceccarelli, G. Vallone, F. De Martini, P. Mataloni, and A. Cabello, Phys. Rev. Lett. **103**, 160401 (2009).
[28] Y.-B. Sheng, F.-G. Deng, and G. L. Long, Phys. Rev. A **82**, 032318 (2010).
[29] B.-C. Ren, H.-R. Wei, M. Hua, T. Li, and F.-G. Deng, arXiv:1207.0168.